

GINGIVAL CREVICULAR FLUID FLOW (VOLUME) AS A BIOMARKER OF ORTHODONTIC TOOTH MOVEMENT

HARRISON TRUMAN¹, NINA STEVENS², MAHBOUBE HASHEMINASAB³, SOLALEH AMIRPOUR³, TYLER STARLEY¹, MARIE M. TOLAROVA⁴, MIROSLAV TOLAR^{4,5}

¹Dental Student; ²Department of Orthodontics Resident; ³Research Fellow; ⁴Department of Orthodontics; ⁵Department of Biomedical Sciences
University of the Pacific Arthur A. Dugoni School of Dentistry, San Francisco, CA, USA



OBJECTIVE

Orthodontic tooth movement (OTM) leads to remodeling of the periodontal ligament, alveolar bone, and gingiva. Tooth movement is characterized by bone deposition at sites of tension and bone resorption at pressure sites (Fig. 1). Due to a sterile inflammatory process in the periodontal ligament (PDL) space, a flow rate of gingival crevicular fluid (GCF) is increased and its composition is modified (Fig. 2). Various cell signaling pathways are activated, which ultimately stimulate PDL turnover, as well as localized bone resorption and bone deposition (Fig. 3). Processes associated with tooth movement may also be better understood through the study of the GCF flow/volume formed in response to such movement. This information could be used clinically to choose a proper mechanical loading, to improve and to shorten a period of treatment, or to avoid adverse consequences, such as root resorption. Our goal is to monitor changes of GCF flow during orthodontic treatment with Invisalign.




Figure 1. Coronal section through the root of a premolar being moved to the left (arrow). This image shows reactions of tissues in PDL space to application of force. The tension zone (A) is wide, strings of osteoblasts are visible reflecting proliferation and migration of osteoblasts, blood vessels are enlarged, the compression zone (B) is narrow, with packed cells and hardly visible blood vessels, lacunae formed by osteoclasts are visible in the alveolar bone, even lacunae in the root surface are seen (C). (Proffit et al. 2018)




Figure 2. Inflammatory response of PDL and innate immune system to movement of a tooth. (Andrade, et al., 2012)




Figure 3. Diagram of bone remodeling. (1) bone that is not remodeled, (2 on the left side) monocytes are attracted into the active remodeling zone and differentiate into osteoclasts, (3) local macrophages digest bone osteoid and release growth factors that were embedded in it, (4) pericytes are the source of new osteoblasts that are triggered by growth factors to proliferate, differentiate and form a new osteoid (2 on the right side). (Matsumoto et al. 2016).

METHODS

Various original studies and systematic reviews were focused on identification of GCF biomarkers and their roles during orthodontic treatment. GCF has been extensively studied because of the simple, quick, noninvasive nature of its collection and ease of repetitive sampling from the same site with the help of Periopaper strips (Oraflow) or micropipettes. GCF is analyzed for the content of various biochemical markers and its volume. We collected GCF (IRB #2021-61) into a Periopaper strip positioned in gingival sulcus for 20 seconds (Fig. 4) and measured its volume using Periotron 8010 (Oraflow).



Figure 4 : Periopaper strip in gingival sulcus

RESULTS

Graphs represent the GCF volumes collected from three subjects (Figs. 5, 6, 7). The curves were grouped in respect to different types of teeth involved. Three general trends can be seen: the curves for teeth of the same type seem to have a similar shape, the curves for one type of tooth seem to differ from the curves of another type, and those curves differed between the three patients.




Figure 5. GCF volume curves of premolars (4, 12, 13, 20, 29), canines (6, 11, 22) and incisors (24, 26) of patient BM21. Timepoints t0 (baseline), t1-t2 first aligner, t3-t4 second aligner, t5-t6 third aligner.




Figure 6. GCF volume curves of premolars (5, 12, 21, 28), canines (6, 11, 22) and incisors (10, 25, 26) of patient BM22. Timepoints 1 (baseline), 2-3 first aligner, 4-5 second aligner, 6-7 third aligner.




Figure 7. GCF volume curves of premolars (12, 28, 29), canines (6, 11, 22) and incisors (7, 9, 23, 25) of patient BM23. Timepoints 1 (baseline), 2-3 first aligner, 4-5 second aligner, 6-7 third aligner.

CONCLUSION

Real-time studies of ongoing processes in periodontal tissues during orthodontic treatment can lead to a proper choice of mechanical loading, shortening of treatment, better planning and minimizing adverse consequences. Our results seem to suggest that GCF volume could be used as a potential biomarker for such decisions. However, future research is needed to quantify tooth movement in three-dimensional PDL space, standardize this movement across teeth of different sizes, and more accurately quantify response of periodontal biomarkers in relation to orthodontic force applied during the treatment.

REFERENCES

- Alansari S, Sangsuwon C, Vongthongleur T, Kwal R, Teo MC, Lee YB. Biological principles behind orthodontic tooth movement. *Semin Orthod.* 2015;21(3):151-161.
- Alhadlaq AM. Biomarkers of Orthodontic Tooth Movement in Gingival Crevicular Fluid: A Systematic Review. *J Contemp Dent Pract* 2015;16(7):578-587.
- Barbieri G, Solano P, Alarcón JA, Vernal R, Rios-Lugo J, Sanz M, et al. Biochemical markers of bone metabolism in gingival crevicular fluid during early orthodontic tooth movement. *The Angle Orthodontist.* 2013;83(1):63-69.
- Brooks PJ, Heckler AF, Wei K, Gong SG. M-CSF accelerates orthodontic tooth movement by targeting preosteoclasts in mice. *Angle Orthod.* 2011;81:277-83.
- d'Apuzzo F, Cappabianca S, Ciavarella D, Monsurrò A, Silvestrini-Biavati A, Perillo L. Biomarkers of periodontal tissue remodeling during orthodontic tooth movement in mice and men: overview and clinical relevance. *ScientificWorldJournal.* 2013 Apr 23;2013:105873. doi: 10.1155/2013/105873. PMID: 23737704; PMCID: PMC3655650.
- Kawasaki K, Takahashi T, Yamaguchi M, Kasai K. Effects of Aging on RANKL and OPG Levels in Gingival Crevicular Fluid During Orthodontic Tooth Movement. *Orthodontics and Craniofacial Research.* 2006;9(3):137-142.
- Krishnan V, Davidovitch Z. Cellular, molecular, and tissue-level reaction to orthodontic force. *Am J Orthod Dentofacial Orthop.* 2006;129(4):469.e1-32.
- Kumar AA, Saravanan K, Kohila K, Kumar SS. Biomarkers in orthodontic tooth movement. *J Pharm Bioall Sci.* 2015;7(6):325-330.
- Li Y, Jacox LA, Little SH, Ko CC. Orthodontic tooth movement: The biology and clinical implications. *Kaohsiung J Med Sci.* 2018 Apr;34(4):207-214. doi: 10.1016/j.kjms.2018.01.007. Epub 2018 Feb 3. PMID: 29655409.
- Matsumoto, Mariza & Bigueti, Claudia & Fonseca, Angélica & Saraiva, Patricia. (2016). Bone Tissue Healing Dynamics: From damage to reconstruction.
- Palmqvist P, Lundberg P, Persson E, Johansson A, Lundgren I, Lie A, et al. Inhibition of hormone and cytokine-stimulated osteoclastogenesis and bone resorption by interleukin-4 and interleukin-13 is associated with increased osteoprotegerin and decreased RANKL and RANK in a STAT6-dependent pathway. *Journal of Biological Chemistry.* 2006;281(5):2414-29.
- Proffit WR, Field HW, Sarver DM. (2018). *Contemporary orthodontics*. St. Louis, Mo: Mosby Elsevier
- Taddei SR, Queiroz-Junior CM, Moura AP, et al. The effect of CCL3 and CCR1 in bone remodeling induced by mechanical loading during orthodontic tooth movement in mice. *Bone.* 2013;52(1):259-267.
- Taddei SR, Andrade I, Queiroz-Junior CM, Garlet TP, Cunha FDQ. Role of CCR2 in orthodontic tooth movement. *American Journal of Orthodontics and Dentofacial Orthopedics.* 2012;44(2):153-160.
- Teixeira CC, Khoo E, Tran J, et al. Cytokine Expression and Accelerated Tooth Movement. *Journal of Dental Research.* 2010;89(10):1135-1141. d

This study was supported by International Award from Align Technology, Inc. We have no financial interest.

OKU Sutro Excellence Day Project Cover Sheet

Project Title: Gingival Crevicular Flow (Volume) as a Biomarker of Orthodontic Tooth Movement

Award Category: DDS & IDS – Research

Contributors to this Project:

1. Harrison Truman, #989395943, DDS 2024
2. Tyler Starley, #989396029, DDS 2024
3. Nina Stevens, #989135186, Ortho Residency 2023
4. Mahboub Hasheminasab, Research Fellow
5. Solaleh Amirpour, Research Fellow
6. Miroslav Tolar, Ortho Faculty
7. Marie Tolarova, Ortho Faculty

Abstract:

Objectives: Tooth movement is caused by the application of force. In short, applied force strains structures present in the PDL space – cells, ligaments, blood vessels. Cells in the PDL are damaged by extension and by diminished oxygen supply due to compression of blood vessels. Compounds released from damaged or dead cells trigger innate inflammatory response. One of the biomarkers of that response is increased formation of extracellular fluid (edema), specifically gingival crevicular fluid (GCF). Our goal is to monitor changes of GCF flow during the orthodontic treatment with Invisalign.

Methods: GCF is collected using Periostrips (Oraflow) from the buccal and lingual sides of maxillary and mandibular teeth (IRB # 2021-61). The volume of GCF is measured using Periotron 8010 (Oraflow). Time points cover the baseline, start, and end of each aligner during the treatment.

Results: GCF volume time series form curves that are typical for each type of tooth but differ in different patients. In literature, it was argued that GCF volume was not a reliable quantitative indicator in respect to tooth movement caused by fixed appliances. We are showing that it is not true for orthodontic tooth movement caused by Invisalign aligners. A possible explanation is discussed.

Conclusions: GCF volume (flow) seems to be a good biomarker of orthodontic tooth movement by Invisalign.

Acknowledgements: This study was supported by Research Award from Align Technology.